
DARIEN DEWOLF, St. Francis Xavier University
Groupoids Associated to Join Inverse Categories

Join inverse categories [1] are inverse categories that come with a way to glue together two partially defined morphisms, provided they agree everywhere that they are defined. Every inverse semigroup can be associated to an inductive groupoid (Ehresmann-Schein-Nambooripad), and by extension every inverse category \mathbf{X} to a top-heavy locally inductive groupoid $\mathcal{G}(\mathbf{X})$ [2].

This talk will show that the groupoids associated to join inverse categories share at least two interesting properties [3]:

(1) Each admits a pair of functors

$$(-)_* : \mathcal{G}(\mathbf{X})^{\text{op}} \rightarrow \mathbf{Loc} \text{ and } (-)^* : \mathcal{G}(\mathbf{X}) \rightarrow \mathbf{Loc}$$

such that, for each arrow $(\alpha : A \rightarrow B) \in \mathcal{G}(\mathbf{X})$, the locale homomorphisms α_* and α^* form an equivalence of categories between A^* and B^* .

(2) Each admits an Ehresmann topology, a data structure very much analogous to a Grothendieck topology, but in the language of *covering order ideals*.

References

- [1] J.R.B. Cockett, G.S.H. Cruttwell, and J.D. Gallagher. Differential restriction categories. *Theory and Applications of Categories*, 25(21):537-613, 2011.
- [2] D. DeWolf and D. Pronk. The Ehresmann-Schein-Nambooripad Theorem for Inverse Categories. arXiv, Nov. 2017, 1507.08615v2.
- [3] D. DeWolf. *Restriction Category Perspectives of Partial Computation and Geometry*. PhD thesis, Dalhousie University, 2017.