
MARIA GRAZIA VIOLA, Lakehead University

Regularities properties of Cuntz-Pimsner algebras associated to C^ -correspondences over commutative C^* -algebras*

We discuss the structural properties of Cuntz-Pimsner algebras arising from full, minimal, non-periodic, and finitely generated projective C^* -correspondence over commutative C^* -algebras. A large class of examples is obtained considering the set $\Gamma(V, \alpha)$ of continuous sections of a complex vector bundle on a compact metric space X , where left multiplication is given by a twist by a minimal homeomorphism $\alpha: X \rightarrow X$.

Cuntz-Pimsner algebras are generalization of both Cuntz-Krieger algebras and crossed products by the integers. In the case of crossed products by minimal homeomorphisms, the orbit breaking subalgebra, defined by I. Putnam, is a large subalgebra in the sense of N. C. Phillips. We show that the Cuntz-Pimsner algebra $\mathcal{O}(\Gamma(V, \alpha))$ also contains a large subalgebras, at least for a large class of C^* -correspondences. We will discuss some properties that $\mathcal{O}(\Gamma(V, \alpha))$ and/or its large subalgebra have, focusing on properties needed for classification by the Elliott invariant.

This is joint work with M. S. Adamo, D. Archey, M. Forough, M. Georgescu, J. A. Jeong, and K. Strung.