

THE OLYMPIAD CORNER

No. 342

Carmen Bruni

Les problèmes présentés dans cette section ont déjà été présentés dans le cadre d'une olympiade mathématique régionale ou nationale. Nous invitons les lecteurs à présenter leurs solutions, commentaires et généralisations pour n'importe quel problème. S'il vous plaît vous référer aux règles de soumission à l'endos de la couverture ou en ligne.

*Pour faciliter l'examen des solutions, nous demandons aux lecteurs de les faire parvenir au rédacteur au plus tard le **1er janvier 2017**; toutefois, les solutions reçues après cette date seront aussi examinées jusqu'au moment de la publication.*

La rédaction souhaite remercier André Ladouceur, Ottawa, d'avoir traduit les problèmes.

OC276. Soit m et n des entiers strictement positifs. Sachant que le nombre

$$k = \frac{(m+n)^2}{4m(m-n)^2 + 4}$$

est un entier, démontrer que k est un carré parfait.

OC277. Déterminer toutes les fonctions $f : \mathbb{R} \rightarrow \mathbb{R}$ qui vérifient

$$f(x^2 + yf(x)) = xf(x + y)$$

pour tous réels x, y .

OC278. Soit n un entier supérieur ou égal à 4. Déterminer toutes les permutations $\{x_1, x_2, \dots, x_n\}$ de $\{1, 2, \dots, n\}$ pour lesquelles $x_i < x_{i+2}$ lorsque $1 \leq i \leq n-2$ et $x_i < x_{i+3}$ lorsque $1 \leq i \leq n-3$.

OC279. Soit ABC un triangle acutangle et O le centre de son cercle circonscrit. Soit P et Q des points sur les côtés respectifs AB et AC tels que

$$BP \cdot CQ = AP \cdot AQ.$$

Soit I un cercle dont le centre est situé sur la hauteur du triangle ABC abaissée au point A et qui passe aux points A, P et Q . Démontrer que I est tangent au cercle circonscrit au triangle BOC .

OC280. Soit $g(n)$ le plus grand commun diviseur de n et 2015. Déterminer le nombre de triplets (a, b, c) qui satisfont aux deux conditions suivantes:

1. $a, b, c \in \{1, 2, \dots, 2015\}$ et
2. $g(a), g(b), g(c), g(a+b), g(b+c), g(c+a), g(a+b+c)$ sont distincts.

.....

OC276. Let m and n be positive integers. If the number

$$k = \frac{(m+n)^2}{4m(m-n)^2 + 4}$$

is an integer, prove that k is a perfect square.

OC277. Find all real functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $f(x^2 + yf(x)) = xf(x+y)$.

OC278. Find all possible permutations $\{x_1, x_2, \dots, x_n\}$ of $\{1, 2, \dots, n\}$ so that when $1 \leq i \leq n-2$ then we have $x_i < x_{i+2}$ and when $1 \leq i \leq n-3$ then we have $x_i < x_{i+3}$. Here $n \geq 4$.


OC279. Let ABC be an acute-angled triangle with circumcenter O . Let I be a circle with centre on the altitude from A in ABC , passing through vertex A and points P and Q on sides AB and AC . Assume that

$$BP \cdot CQ = AP \cdot AQ.$$

Prove that I is tangent to the circumcircle of triangle BOC .


OC280. Let $g(n)$ be the greatest common divisor of n and 2015. Find the number of triples (a, b, c) which satisfy the following two conditions:

1. $a, b, c \in 1, 2, \dots, 2015$;
2. $g(a), g(b), g(c), g(a+b), g(b+c), g(c+a), g(a+b+c)$ are pairwise distinct.



OLYMPIAD SOLUTIONS

Les énoncés des problèmes dans cette section paraissent initialement dans 2015: 41(2), p. 55–56.



OC216. Let $p = n^2 + 1$ be a given prime number. Find the set of integer solutions to the equation below:

$$x^2 - (n^2 + 1)y^2 = n^2.$$

Originally problem 4 from the number theory portion of the third round of the 2013 Iranian National Mathematical Olympiad.

We received 2 correct submissions. We present the solution by Oliver Geupel.

We will show that the solutions for $p = 2$ are $x + y\sqrt{2} = \pm(3 + 2\sqrt{2})^k$, where $k \in \mathbb{Z}$. We will prove that the solutions for $p > 2$ are

$$\begin{aligned} x + y\sqrt{p} \in \{ & \pm (2n^2 + 1 + 2n\sqrt{p})^k n, \\ & \pm (2n^2 + 1 + 2n\sqrt{p})^k (n^2 - n + 1 + (n - 1)\sqrt{p}), \\ & \pm (2n^2 + 1 + 2n\sqrt{p})^k (n^2 + n + 1 + (n + 1)\sqrt{p}) \mid k \in \mathbb{Z} \}. \end{aligned}$$

For $z = x + y\sqrt{p} \in \mathbb{Z}[\sqrt{p}]$, denote $N(z) = x^2 - py^2$. We recapitulate the following well-known facts on Pell-like equations (check, e.g., the article *Pell's Equations* by Dušan Dukić on the website of the IMO compendium, retrieved March 25, 2016 from <http://imomath.com/index.php?option=615>):

- (1) If z_1 is the fundamental solution (which must exist) of the equation $N(z) = 1$, i.e., the minimal element of $\mathbb{Z}[\sqrt{p}]$ with $z > 1$ and $N(z) = 1$, then all the solutions $z \in \mathbb{Z}[\sqrt{p}]$ are given by $z = \pm z_1^k$, $k \in \mathbb{Z}$.
- (2) The fundamental solution of the equation $x^2 - 2y^2 = 1$ is $3 + 2\sqrt{2}$.
- (3) For $a \in \mathbb{Z}$, every solution of the equation $N(z) = a$ has the form $z = \pm z_1^k z_a$ ($k \in \mathbb{Z}$) where z_1 is the fundamental solution of the equation $N(z) = 1$, and $z_a = x_a + y_a\sqrt{p}$ is a solution of $N(z) = a$ with $1 \leq z_a \leq z_1$. Also

$$|x_a| \leq \frac{z_1 + 1}{2\sqrt{z_1}} \sqrt{|a|}.$$

The result for $p = 2$ follows from (1) and (2). It remains to consider $p > 2$.

Let $z_1 = x_1 + y_1\sqrt{p}$ be the fundamental solution of the equation $N(z) = 1$. We obtain $y_1^2 = (x_1 - 1)(x_1 + 1)/p$, that is, p is a divisor of either $x_1 - 1$ or $x_1 + 1$. We check the small values for x_1 in succession. If $x_1 = p - 1$ then $y_1^2 + 1 = n^2$, which is impossible. If $x_1 = p + 1$ then $y_1^2 = n^2 + 3$ which is impossible for $n > 1$. Trying $x_1 = 2p - 1$, we obtain $z_1 = 2n^2 - 1 + 2n\sqrt{p}$.

To complete the work, by (3) it is enough to find the solutions $z_a = x_a + y_a\sqrt{p}$ of $N(z_a) = a$ with $a = n^2$ and


$$x_a < \frac{2n^2 + 2n\sqrt{n^2 + 1}}{2\sqrt{2n^2 - 1} + 2n\sqrt{n^2 + 1}} \cdot n < 2n^2.$$

From $N(z_a) = a$ we have $y_a^2 = (x_a - n)(x_a + n)/p$, that is, p divides either $x_a - n$ or $x_a + n$. We check the small values for x_a in succession. Trying $x_a = n$, we find $z_a = n$. Putting $x_a = p - n$, we obtain $z_a = n^2 - n + 1 + (n - 1)\sqrt{p}$. For $x_a = p + n$, we get $z_a = n^2 + n + 1 + (n + 1)\sqrt{p}$. Trying $x_a = 2p - n$, we obtain $y_a^2 = 4(p - n)$. Hence, $p - n = n^2 + 1 - n$ is a perfect square, say m^2 . We obtain $(n - 1)^2 < m^2 < n^2$, a contradiction. The solution is complete.

OC217. Let G be the centroid of a right-angled triangle ABC with $\angle BCA = 90^\circ$. Let P be the point on ray AG such that $\angle CPA = \angle CAB$, and let Q be the point on ray BG such that $\angle CQB = \angle ABC$. Prove that the circumcircles of triangles AQG and BPG meet at a point on side AB .

Originally problem 3 of the 2013 Canadian Mathematical Olympiad.

We received 3 correct submissions. We present the solution by Andrea Fanchini.



We use barycentric coordinates and the usual Conway's notations with reference to triangle ABC . With $\angle BCA = 90^\circ$, we have $S_C = 0$ so $S_A = b^2$, $S_B = a^2$, $c^2 = a^2 + b^2$. Then a point P on ray AG has coordinates $P(u : 1 : 1)$ where u is a parameter. Now the oriented angle δ (with $0 \leq \delta \leq \pi$) between two lines

$d_i \equiv p_i x + q_i y + r_i z = 0 (i = 1, 2)$, is given from

$$S_\delta = S \cot \delta = \frac{S_A(q_1 - r_1)(q_2 - r_2) + S_B(r_1 - p_1)(r_2 - p_2) + S_C(p_1 - q_1)(p_2 - q_2)}{\begin{vmatrix} 1 & 1 & 1 \\ p_1 & q_1 & r_1 \\ p_2 & q_2 & r_2 \end{vmatrix}}.$$

Therefore using the above, the angle between the line $CP \equiv x - uy = 0$ and the median $AP \equiv y - z = 0$, is

$$S_\delta = \frac{a^2 - 2b^2 u}{u + 2}.$$

But $\angle CPA = \angle CAB$, so

$$S_\delta = S_A = b^2 \Rightarrow u = \frac{a^2 - 2b^2}{3b^2} \Rightarrow P(a^2 - 2b^2 : 3b^2 : 3b^2).$$

Now the equation of a circle is $a^2 yz + b^2 zx + c^2 xy - (x + y + z)(px + qy + rz) = 0$, so to find the equation of the circumcircle of $\triangle BPG$ we have to put in the coordinates of $B(0 : 1 : 0)$, $G(1 : 1 : 1)$, $P(a^2 - 2b^2 : 3b^2 : 3b^2)$ and solving the system, we have $p = b^2, q = 0, r = \frac{2a^2 - b^2}{3}$. Therefore, the intersection between the circumcircle BPG and the side AB gives the point X

$$\begin{cases} a^2 yz + b^2 zx + c^2 xy - (x + y + z)(b^2 x + \frac{2a^2 - b^2}{3} z) = 0 \\ z = 0 \end{cases} \Rightarrow X(a^2 : b^2 : 0).$$

In the same way, we have $Q(1 : v : 1)$ and the angle between the median $BQ \equiv x - z = 0$ and the line $CQ \equiv vx - y = 0$, is

$$S_\epsilon = \frac{b^2 - 2a^2 v}{v + 2}.$$

But $\angle CQB = \angle ABC$, so

$$S_\epsilon = S_B = a^2 \Rightarrow v = \frac{b^2 - 2a^2}{3a^2} \Rightarrow Q(3a^2 : b^2 - 2a^2 : 3a^2).$$

To find the equation of the circumcircle of $\triangle AQQ$, we have to put in the coordinates of $A(1 : 0 : 0)$, $G(1 : 1 : 1)$, $Q(3a^2 : b^2 - 2a^2 : 3a^2)$ and solving the system, we have $p = 0, q = a^2, r = \frac{2b^2 - a^2}{3}$. Therefore, the intersection between the circumcircle AQG and the side AB gives the point X'

$$\begin{cases} a^2 yz + b^2 zx + c^2 xy - (x + y + z)(a^2 y + \frac{2b^2 - a^2}{3} z) = 0 \\ z = 0 \end{cases} \Rightarrow X'(a^2 : b^2 : 0).$$

So we have $X \equiv X'$, as required.

Note: this point is the intersection of the symmedian through vertex C and the side AB . The three symmedians concur at the Symmedian Point $K(a^2 : b^2 : c^2)$, well known also as Lemoine point.

OC218. Find all functions $f : \mathbb{N} \rightarrow \mathbb{N}$ satisfying

$$f(mn) = \text{lcm}(m, n) \cdot \gcd(f(m), f(n))$$

for all positive integers m, n .

Originally problem 5 from day 2 of the 2013 Korean National Olympiad.

We have a partial solution submitted by Konstantine Zelator which we include (and complete) here.

We claim that the only solutions are $f(m) = km$ for any $k \in \mathbb{N}$. First, this function is a solution since

$$\text{lcm}(m, n) \cdot \gcd(f(m), f(n)) = \frac{mn}{\gcd(m, n)} \cdot \gcd(km, kn) = kmn = f(mn).$$

Now, let's call the given relation $R_{m,n}$ and suppose $f(1) = c$. Then, $R_{m,1}$ for any $m \in \mathbb{N}$ gives

$$f(m) = m \gcd(f(m), c).$$

Now, by the relation $R_{cm,1}$ for any $m \in \mathbb{N}$ (and using the above), we see that

$$f(cm) = cm \cdot \gcd(f(cm), c) = cm \cdot \gcd(c \cdot \gcd(f(m), c), c) = c^2 m.$$

Next, the relation $R_{m,cn}$ for any $m, n \in \mathbb{N}$ reveals with the above two facts that

$$c^2 mn = f(mcn) = \text{lcm}(m, cn) \cdot \gcd(f(m), f(cn)) = \frac{cmn}{\gcd(m, cn)} \cdot \gcd(f(m), c^2 n)$$

and simplifying yields

$$c \gcd(m, cn) = \gcd(f(m), c^2 n).$$

Therefore, $c \mid f(m)$. Hence, from the first derived relation above, we see that

$$f(m) = m \gcd(f(m), c) = mc$$

completing the proof.

OC219. Given positive integers m and n , prove that there is a positive integer c such that the numbers cm and cn have the same number of occurrences of each non-zero digit when written in base ten.

Originally problem 5 from day 2 of the 2013 USAMO.

We present the solution by Oliver Geupel. There were no other submissions.

Without loss of generality assume $m < n$. We consider the cases $\gcd(m, 10) = 1$ and $\gcd(m, 10) > 1$ in succession.

First, consider the case $\gcd(m, 10) = 1$. We have $\gcd(10^{n^2}n - m, 10) = 1$. Hence, there exists an integer $a > 3n^2$ such that $10^a \equiv 1 \pmod{10^{n^2}n - m}$. Fix such

an integer a and put $b = a - 2n^2$. As a consequence, we have $b > n^2$. Note that $m \equiv 10^{n^2}n \pmod{10^{n^2}n - m}$. Hence,

$$10^{n^2+b}m \equiv 10^{2n^2+b}n \equiv n \pmod{10^{n^2}n - m},$$

that is, the number $d = \frac{10^{n^2+b}m - n}{10^{n^2}n - m}$ is an integer.

We are going to show that

$$c = 10^{3b} + d$$

has the desired property.

By direct computation, we have

$$cm = 10^{3b}m + 10^{n^2} \cdot \frac{10^b m^2 - n^2}{10^{n^2}n - m} + n, \quad cn = 10^{3b}n + 10^b m + \frac{10^b m^2 - n^2}{10^{n^2}n - m}.$$

Thus,

$$f = \frac{10^b m^2 - n^2}{10^{n^2}n - m}$$

is an integer. We have $0 < f < 10^b$. Consequently, the decimal expansion of cm is the concatenation of the decimal expansions of the three numbers m , f , and n , padded with some additional zeros. Also, the decimal representation of cn is the concatenation of the decimal representations of n , m , and f , padded with some extra zeros. This completes the proof in the case $\gcd(m, 10) = 1$.

It remains to examine the case $\gcd(m, 10) > 1$. Let $m = 2^h 5^j k$ with nonnegative integers h , j , and k where $\gcd(k, 10) = 1$. For positive integers x and y , we write $x \sim y$ if the decimal representations of x and y have the same number of occurrences of each nonzero digit. We know that there is a positive integer c_0 such that $c_0 k \sim c_0 2^j 5^h n$. We obtain

$$2^j 5^h c_0 m = c_0 10^{h+j} k \sim c_0 k \sim 2^j 5^h c_0 n.$$

Consequently, the number

$$c = 2^j 5^h c_0$$

has the required property for m and n .

OC220. Let $A_1 A_2 \dots A_8$ be a convex octagon such that all of its sides are equal and its opposite sides are parallel. For each $i = 1, \dots, 8$, define B_i as the intersection between segments $A_i A_{i+4}$ and $A_{i-1} A_{i+1}$, where $A_{j+8} = A_j$ and $B_{j+8} = B_j$ for all j . Show some number i , amongst 1, 2, 3, and 4 satisfies

$$\frac{A_i A_{i+4}}{B_i B_{i+4}} \leq \frac{3}{2}.$$

Originally problem 6 of the 2013 Mexican National Olympiad.

No submitted solutions.